

Name: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_ MR: \_\_\_\_\_

## PATIENT HEALTH QUESTIONNAIRE-9 (PHQ-9)

Over the last 2 weeks, how often have you been bothered  
by any of the following problems?

(Use "✓" to indicate your answer)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3

FOR OFFICE CODING 0 + \_\_\_\_\_ + \_\_\_\_\_ + \_\_\_\_\_

=Total Score: \_\_\_\_\_

If you checked off any problems, how difficult have these problems made it for you to do your  
work, take care of things at home, or get along with other people?

Not difficult at all <input type="checkbox"/>	Somewhat difficult <input type="checkbox"/>	Very difficult <input type="checkbox"/>	Extremely difficult <input type="checkbox"/>
---	---	---	--

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from  
Pfizer Inc. No permission required to reproduce, translate, display or distribute.

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

## GAD-7

Over the last 2 weeks, how often have you been bothered by the following problems?

Not  
at all

Several  
days

More than  
half the  
days

Nearly  
every day

(Use "✓" to indicate your answer)

1. Feeling nervous, anxious or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it is hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid as if something awful might happen	0	1	2	3

(For office coding: Total Score T \_\_\_\_ = \_\_\_\_ + \_\_\_\_ + \_\_\_\_)

Severity Range:

\_\_\_ 0-4: Minimum

\_\_\_ 5-9: Mild

\_\_\_ 10-14: Moderate

\_\_\_ 15-21: Severe

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

**COLUMBIA-SUICIDE SEVERITY RATING SCALE**  
 **Screener/Recent – Self-Report**

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

		In The Past Month	
Answer Questions 1 and 2		YES	NO
1) <i>Have you wished you were dead or wished you could go to sleep and not wake up?</i>			
2) <i>Have you actually had any thoughts about killing yourself?</i>			
If YES to 2, answer questions 3, 4, 5, and 6. If NO to 2, go directly to question 6			
3) <i>Have you thought about how you might do this?</i>			
4) <i>Have you had any intention of acting on these thoughts of killing yourself, as opposed to you have the thoughts but you definitely would not act on them?</i>			
5) <i>Have you started to work out or worked out the details of how to kill yourself?</i> <i>Do you intend to carry out this plan?</i>			
		In the Past 3 Months	
6) <i>Have you done anything, started to do anything, or prepared to do anything to end your life?</i>  Examples: Collected pills, obtained a gun, gave away valuables, wrote a will or suicide note, took out pills but didn't swallow any, held a gun but changed your mind or it was grabbed from your hand, went to the roof but didn't jump; or actually took pills, tried to shoot yourself, cut yourself, tried to hang yourself, etc.			
<i>In your entire lifetime, how many times have you done any of these things?</i>			

Therapist Name: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_



Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

### **Alcohol Use Disorders Identification Test – AUDIT**

*Please select the answer that is most correct for you to each of the following questions.*

- 1) How often do you have a drink containing alcohol? *(If you answer never, jump to questions 9&10)*  
(0) Never (1) Monthly or less (2) 2-4 times a month (3) 2-3 times per week (4) 4 or more times a week
- 2) How many drinks containing alcohol do you have on a typical day when you are drinking?  
(0) 1-2 (1) 3-4 (2) 5-6 (3) 7-8 (4) 10 or more
- 3) How often do you have six or more drinks on one occasion?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 4) How often during the last year have you found that you were not able to stop drinking once you had started?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 5) How often during the last year have you failed to do what was normally expected from you because of drinking?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 6) How often during the last year have you needed a first drink in the morning to get yourself going after a heavy drinking session?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 7) How often during the last year have you had a feeling of guilt or remorse after drinking?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 8) How often during the last year have you been unable to remember what happened the night before because you had been drinking?  
(0) Never (1) Less than monthly (2) Monthly (3) Weekly (4) Daily or almost daily
- 9) Have you or someone else been injured as a result of your drinking?  
(0) No (2) Yes, but not in the last year (3) Yes, during the last year
- 10) Has a relative or friend, or a doctor or other health worker, been concerned about your drinking or suggested you cut down?  
(0) No (2) Yes, but not in the last year (3) Yes, during the last year

Total Score: \_\_\_\_\_

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

MR: \_\_\_\_\_

Patient's Name: \_\_\_\_\_

Date: \_\_\_\_\_

**Drug Abuse Screening Test—DAST-10**

These Questions Refer to the Past 12 Months

1	Have you used drugs other than those required for medical reasons?	Yes	No
2	Do you abuse more than one drug at a time?	Yes	No
3	Are you unable to stop using drugs when you want to?	Yes	No
4	Have you ever had blackouts or flashbacks as a result of drug use?	Yes	No
5	Do you ever feel bad or guilty about your drug use?	Yes	No
6	Does your spouse (or parents) ever complain about your involvement with drugs?	Yes	No
7	Have you neglected your family because of your use of drugs?	Yes	No
8	Have you engaged in illegal activities in order to obtain drugs?	Yes	No
9	Have you ever experienced withdrawal symptoms (felt sick) when you stopped taking drugs?	Yes	No
10	Have you had medical problems as a result of your drug use (eg, memory loss, hepatitis, convulsions, bleeding)?	Yes	No

Total Score: \_\_\_\_\_

**Guidelines for Interpretation of DAST-10**

Interpretation (Each "Yes" response = 1)

Score	Degree of Problems Related to Drug Abuse	Suggested Action
0	No problems reported	Encouragement and education
1-2	Low level	Risky behavior – feedback and advice
3-5	Moderate level	Harmful behavior – feedback and counseling; possible referral for specialized assessment
6-8	Substantial level	Intensive assessment and referral

Skinner HA. The Drug Abuse Screening Test. *Addictive Behavior*. 1982;7(4):363-371.

Yudko E, Lozhkina O, Fouts A. A comprehensive review of the psychometric properties of the Drug Abuse Screening Test.

*J Subst Abuse Treatment* 2007;32:189-198

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_



Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

### **Rosenberg's Self-Esteem Scale** (Rosenberg, 1965)

**Instructions:** Below is a list of statements dealing with your general feelings about yourself. Please indicate how strongly you agree or disagree with each statement.

	<b>Strongly Agree</b>	<b>Agree</b>	<b>Disagree</b>	<b>Strongly Disagree</b>
1. I feel that I'm a person of worth, at least on an equal plane with others.				
2. I feel that I have a number of good qualities.				
3. All in all, I am inclined to feel that I am a failure.				
4. I am able to do things as well as most other people.				
5. I feel I do not have much to be proud of.				
6. I take a positive attitude toward myself.				
7. On the whole, I am satisfied with myself.				
8. I wish I could have more respect for myself.				
9. I certainly feel useless at times.				
10. At times I think I am no good at all.				

Total Score: \_\_\_\_\_

**Scoring:** To score the items, assign a value to each of the 10 items as follows:

- **For items 1, 2, 4, 6, 7:** Strongly Agree=3, Agree=2, Disagree=1, and Strongly Disagree=0.
- **For items 3, 5, 8, 9, 10** (which are reversed in valence, and noted with the asterisks\*\* below): Strongly Agree=0, Agree=1, Disagree=2, and Strongly Disagree=3.

The scale ranges from 0-30, with 30 indicating the highest score possible. Higher scores indicate higher self-esteem. Scores below 15 indicate low self-esteem.

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Name: \_\_\_\_\_ MR: \_\_\_\_\_ Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

### Satisfaction With Life Scale (SWLS)

Instructions: Below are five statements that you may agree or disagree with. Using the 1 - 7 scale below, indicate your agreement with each item by placing the appropriate number on the line preceding that item. Please be open and honest in your responding.

7 - Strongly agree

6 - Agree

5 - Slightly agree

4 - Neither agree nor disagree

3 - Slightly disagree

2 - Disagree

1 - Strongly disagree

\_\_\_\_ In most ways my life is close to my ideal.

\_\_\_\_ The conditions of my life are excellent.

\_\_\_\_ I am satisfied with my life.

\_\_\_\_ So far I have gotten the important things I want in life.

\_\_\_\_ If I could live my life over, I would change almost nothing.

Total Score: \_\_\_\_\_

Severity Range:

31 – 35: Extremely satisfied

26 – 30: Satisfied

21 – 25: Slightly satisfied

20: Neutral

15 – 19: Slightly dissatisfied

10 – 14: Dissatisfied

5 - 9: Extremely dissatisfied

Ed Diener, Robert A. Emmons, Randy J. Larsen and Sharon Griffin as noted in the 1985 article in the *Journal of Personality Assessment*.

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

Name of Caregiver: \_\_\_\_\_ Relationship: \_\_\_\_\_

Patient Name: \_\_\_\_\_ Date: \_\_\_\_\_

Patient ID # \_\_\_\_\_

## Katz Index of Independence in Activities of Daily Living

Activities Points (1 or 0)	Independence (1 Point)	Dependence (0 Points)
	<b>NO</b> supervision, direction or personal assistance.	<b>WITH</b> supervision, direction, personal assistance or total care.
<b>BATHING</b> Points: _____	<b>(1 POINT)</b> Bathes self completely or needs help in bathing only a single part of the body such as the back, genital area or disabled extremity.	<b>(0 POINTS)</b> Need help with bathing more than one part of the body, getting in or out of the tub or shower. Requires total bathing
<b>DRESSING</b> Points: _____	<b>(1 POINT)</b> Get clothes from closets and drawers and puts on clothes and outer garments complete with fasteners. May have help tying shoes.	<b>(0 POINTS)</b> Needs help with dressing self or needs to be completely dressed.
<b>TOILETING</b> Points: _____	<b>(1 POINT)</b> Goes to toilet, gets on and off, arranges clothes, cleans genital area without help.	<b>(0 POINTS)</b> Needs help transferring to the toilet, cleaning self or uses bedpan or commode.
<b>TRANSFERRING</b> Points: _____	<b>(1 POINT)</b> Moves in and out of bed or chair unassisted. Mechanical transfer aids are acceptable	<b>(0 POINTS)</b> Needs help in moving from bed to chair or requires a complete transfer.
<b>CONTINENCE</b> Points: _____	<b>(1 POINT)</b> Exercises complete self control over urination and defecation.	<b>(0 POINTS)</b> Is partially or totally incontinent of bowel or bladder
<b>FEEDING</b> Points: _____	<b>(1 POINT)</b> Gets food from plate into mouth without help. Preparation of food may be done by another person.	<b>(0 POINTS)</b> Needs partial or total help with feeding or requires parenteral feeding.
<b>TOTAL POINTS:</b> _____ <b>SCORING:</b> 6 = High ( <i>patient independent</i> ) 0 = Low ( <i>patient very dependent</i> )		

Source:

try this: Best Practices in Nursing Care to Older Adults, The Hartford Institute for Geriatric Nursing, New York University, College of Nursing, [www.hartfordnig.org](http://www.hartfordnig.org).

Therapist: \_\_\_\_\_ Therapist Signature: MaineHealth



Name of Caregiver: \_\_\_\_\_ Relationship: \_\_\_\_\_

Patient Name: \_\_\_\_\_

Date: \_\_\_\_\_

Patient ID # \_\_\_\_\_

**LAWTON - BRODY  
INSTRUMENTAL ACTIVITIES OF DAILY LIVING SCALE (I.A.D.L.)**

**Scoring:** For each category, circle the item description that most closely resembles the client's highest functional level (either 0 or 1).

<b>A. Ability to Use Telephone</b>		<b>E. Laundry</b>	
1. Operates telephone on own initiative-looks up and dials numbers, etc.	1	1. Does personal laundry completely	1
2. Dials a few well-known numbers	1	2. Launders small items-rinses stockings, etc.	1
3. Answers telephone but does not dial	1	3. All laundry must be done by others	0
4. Does not use telephone at all	0		
<b>B. Shopping</b>		<b>F. Mode of Transportation</b>	
1. Takes care of all shopping needs independently	1	1. Travels independently on public transportation or drives own car	1
2. Shops independently for small purchases	0	2. Arranges own travel via taxi, but does not otherwise use public transportation	1
3. Needs to be accompanied on any shopping trip	0	3. Travels on public transportation when accompanied by another	1
4. Completely unable to shop	0	4. Travel limited to taxi or automobile with assistance of another	0
		5. Does not travel at all	0
<b>C. Food Preparation</b>		<b>G. Responsibility for Own Medications</b>	
1. Plans, prepares and serves adequate meals independently	1	1. Is responsible for taking medication in correct dosages at correct time	1
2. Prepares adequate meals if supplied with ingredients	0	2. Takes responsibility if medication is prepared in advance in separate dosage	0
3. Heats, serves and prepares meals, or prepares meals, or prepares meals but does not maintain adequate diet	0	3. Is not capable of dispensing own medication	0
4. Needs to have meals prepared and served	0		
<b>D. Housekeeping</b>		<b>H. Ability to Handle Finances</b>	
1. Maintains house alone or with occasional assistance (e.g. "heavy work domestic help")	1	1. Manages financial matters independently (budgets, writes checks, pays rent, bills, goes to bank), collects and keeps track of income	1
2. Performs light daily tasks such as dish washing, bed making	1	2. Manages day-to-day purchases, but needs help with banking, major purchases, etc.	1
3. Performs light daily tasks but cannot maintain acceptable level of cleanliness	1	3. Incapable of handling money	0
4. Needs help with all home maintenance tasks	1		
5. Does not participate in any housekeeping tasks	0		
<b>Score</b>		<b>Score</b>	

**Total score** \_\_\_\_\_

A summary score ranges from 0 (low function, dependent) to 8 (high function, independent) for women and 0 through 5 for men to avoid potential gender bias.

Source: *try this*: Best Practices in Nursing Care to Older Adults, The Hartford Institute for Geriatric Nursing, New York University, College of Nursing, [www.hartfordign.org](http://www.hartfordign.org).

Therapist: \_\_\_\_\_ Therapist Signature: \_\_\_\_\_

**MaineHealth**